


THE LOWER G.I. TRACT: ORGANICALLY SOUND



Double-contrast radiography utilizes an air-barium enema to locate or better visualize colonic lesions, and may be of value in selected patients to rule out organic etiology in abdominal dysfunctions.

...BUT OVERSENSITIVE TO EMOTIONAL STRESS

IN IRRITABLE BOWEL SYNDROME* LIBRAX PROVIDES DISTINCTIVE ADVANTAGES

- the specific antianxiety action of LIBRIUM® (chlordiazepoxide HCl)
- the potent antispasmodic action of QUARZAN® (clidinium Br)



Adjunctive
LIBRAX®

Each capsule contains
5 mg chlordiazepoxide HCl
and 2.5 mg clidinium Br.

A clear advantage for patients with
irritable bowel syndrome


ROCHE

*This drug has been evaluated as possibly effective for this indication.
Please see following page for brief summary of prescribing information.

A CLEAR ADVANTAGE FOR PATIENTS WITH IRRITABLE BOWEL SYNDROME*

Adjunctive
LIBRAX®

Each capsule contains 5 mg chlordiazepoxide HCl and 2.5 mg clidinium Br.



The initial prescription allows evaluation of patient response to therapy.

Follow-up therapy with a prescription for a 2- to 3-week supply of medication usually helps maintain patient gains.

ONLY LIBRAX PROVIDES THE SPECIFIC ANTIANXIETY ACTION OF LIBRIUM® (chlordiazepoxide HCl) PLUS THE POTENT ANTISPASMODIC ACTION OF QUARZAN® (clidinium Br)

Please consult complete prescribing information, a summary of which follows:

- * **Indications:** Based on a review of this drug by the National Academy of Sciences—National Research Council and/or other information, FDA has classified the indications as follows:
- "Possibly" effective: as adjunctive therapy in the treatment of peptic ulcer and in the treatment of the irritable bowel syndrome (irritable colon, spastic colon, mucous colitis) and acute enterocolitis.
- Final classification of the less-than-effective indications requires further investigation.

Contraindications: Glaucoma; prostatic hypertrophy, benign bladder neck obstruction; hypersensitivity to chlordiazepoxide HCl and/or clidinium Br.

Warnings: Caution patients about possible combined effects with alcohol and other CNS depressants, and against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving). Physical and psychological dependence rarely reported on recommended doses, but use caution in administering Librium® (chlordiazepoxide HCl) to known addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions) reported following discontinuation of the drug.

Usage in Pregnancy: Use of minor tranquilizers during first trimester should almost always be avoided because of increased risk of congenital malformations as suggested in several studies. Consider possibility of pregnancy when instituting therapy. Advise patients to discuss therapy if they intend to or do become pregnant.

As with all anticholinergics, inhibition of lactation may occur.

Precautions: In elderly and debilitated, limit dosage to smallest effective amount to preclude ataxia, oversedation, confusion (no more than 2 capsules/day initially; increase gradually as needed and tolerated). Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider pharmacology of agents, particularly potentiating drugs such as MAO in-

hibitors, phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function. Paradoxical reactions reported in psychiatric patients. Employ usual precautions in treating anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship not established.

Adverse Reactions: No side effects or manifestations not seen with either compound alone reported with Librax. When chlordiazepoxide HCl is used alone, drowsiness, ataxia, confusion may occur, especially in elderly and debilitated; avoidable in most cases by proper dosage adjustment, but also occasionally observed at lower dosage ranges. Syncope reported in a few instances. Also encountered: isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent, generally controlled with dosage reduction; changes in EEG patterns may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice, hepatic dysfunction reported occasionally with chlordiazepoxide HCl, making periodic blood counts and liver function tests advisable during protracted therapy. Adverse effects reported with Librax typical of anticholinergic agents, i.e., dryness of mouth, blurring of vision, urinary hesitancy, constipation. Constipation has occurred most often when Librax therapy is combined with other spasmolytics and/or low residue diets.

Dosage: Individualize for maximum benefit. Usual maintenance dose is 1-2 capsules, 3-4 times/day, before meals and at bedtime. Geriatric patients—see Precautions.

How Supplied: Available in green capsules, each containing 5 mg chlordiazepoxide HCl (Librium®) and 2.5 mg clidinium Br (Quarzan®)—bottles of 100 and 500; Tel-E-Dose® packages of 100; Prescription Paks of 50, singly and in trays of 10.



Roche Products Inc.
Manati, Puerto Rico 00701

**BURROUGHS WELLCOME CO. MAKES
CODEINE COMBINATION PRODUCTS.
YOU MAKE THE CHOICE.**



**EMPIRIN[®]
COMPOUND
c̄ CODEINE
#3**

Each tablet contains:
codeine phosphate, 32 mg (gr ½),
(Warning: May be habit-forming);
aspirin, 227 mg; phenacetin, 162 mg;
and caffeine, 32 mg.



**EMPRACET[™]
c̄ CODEINE
#3**

Each tablet contains:
codeine phosphate, 30 mg (gr ½),
(Warning: May be habit-forming);
and acetaminophen 300 mg.



Wellcome

Burroughs Wellcome Co.
Research Triangle Park
North Carolina 27709

What kind of person becomes a Navy physician?



Doctors just like you.

Contrary to what you may think, Navy doctors start their medical careers just like you. In a civilian setting.

In fact, if you look at the personal backgrounds, schooling and medical experience of each and every Navy physician, you'd be hard put to identify a typical Navy physician. Because Navy physicians come from all parts of the country, with a wide range of medical experience. From Park Avenue to Main Street, U.S.A. From interns, to doctors with 20 years' experience. In truth, the Navy physician is you.

But what makes a doctor become a Navy physician?

Lots of reasons.

A minimum of paperwork.

As a Navy physician, we feel your time is too valuable to spend on administrative details. So they're kept to a minimum. Instead, a highly trained staff of professionals attends to the paperwork. The result is that your time can be spent doing what you studied so hard for—practicing medicine.

A challenging practice.

And the practice you'll be given is as varied and challenging as any you'll find in a civilian setting. You'll be treating active duty, dependent and retired personnel—from infant to geriatrics.

Responsibility.

The medical decisions you make will be yours and yours alone. There'll

be no one looking over your shoulder offering second guesses or opinions. But when needed, professional consultation is always available.

Insurance.

The difference between civilian and military insurance is staggering. As a practicing Navy physician, you'll receive free professional liability protection under the Federal Tort Claims Act.

Other benefits.

There are plenty of other great benefits that go with being a Navy physician. Good pay—\$30,000 to start, more depending on your experience. A family life with time for your family. The opportunity to fur-

ther your training. Association with other highly motivated physicians. Even 30 days' paid vacation a year.

The best way to get all the facts is to mail the coupon, or call the Medical Recruiter, toll-free, at **800-841-8000**. (In Georgia, 800-342-5855. In these locations, call collect: Alaska, 272-9133, Puerto Rico, 724-4525.)

You owe it to yourself to get all the facts on a great way to practice medicine.

**Be the doctor
you want to be.
In the Navy.**

.....

• • • • •

• LT. J. D. TAYLOR, MSC, USNR
• Navy Recruiting Area EIGHT
• Building One, NAVSUPACT T.I.
• San Francisco, California 94130
• Tel. (415) 765-5382 (collect)

• (H-1)

• B256

• NAME _____
• First (Please Print or Type) Last

• STREET _____ CITY _____

• STATE _____ ZIP _____ PHONE _____
• (Area Code & No.)

• † MEDICAL SCHOOL _____

• † YEAR GRADUATED _____ DATE OF BIRTH _____

• I AM INTERESTED IN
• (CHECK ONE):

• ☐ Flight Surgeon
• ☐ Undersea Medicine
• ☐ General Medical Officer
• ☐ Practicing My Specialty

• *MY SPECIALTY (IF ANY) IS:

• ☐ Anesthesiology
• ☐ Family Practice
• ☐ Psychiatry
• ☐ Internal Medicine

• ☐ Neurology
• ☐ Radiology
• ☐ Pathology
• ☐ Pediatrics

• STATUS ☐
• (CHECK ONE):

• ☐ Private Practice
• ☐ Hospital Staff
• ☐ Intern
• ☐ Resident

COLBY PROCLAIMS WOMAN SUFFRAGE

Signs Certificate of Ratification
at His Home Without
Women Witnesses.

MILITANTS VEXED AT PRIVACY.

Wanted Movies of Ceremony,
But Both Factions Are

WASHINGTON, Aug. 26, 1920—



Social Security Bill Is Signed; Gives Pensions to Aged, Jobs

Roosevelt Approves Message Intended to Benefit 30,000
Persons When States Adopt Cooperating Laws—He Calls
the Measure 'Cornerstone' of His Economic Program

SENATE APPROVES 18-YEAR OLD VOTE IN ALL ELECTIONS

Amendment to Constitution
is Sent to House, Where
Passage is Expected

WASHINGTON, March 10,
1971—The Senate approved

WASHINGTON, Aug. 14,
The Social Security Bill, part
a broad program of unemployment
insurance and old age pensions
and counted upon to benefit
20,000,000 persons, became law
today when it was signed by
President Roosevelt in the presence
of those chiefly responsible for
bringing it through Congress.

Mr. Roosevelt called it
"the cornerstone
which is being
laid for the
new social order."

TRUMAN CLOSES UNITED NATIONS CONFERENCE WITH PLEA TO TRANSLATE CHARTER INTO DEEDS

NEW WORLD HOPE

President Hails 'Great
Instrument of Peace,'
Insists It Be Used

HISTORIC LANDMARK

Meeting Gives Standing
Ovation as Executive
Promises Peace Gain

SAN FRANCISCO, June 26, 1945

"If we fail to use it," he declared
to the solemn final meeting of the
delegates, 'we shall betray all of
those who have died in order that
we might meet here in freedom and
safety to create it.'

"If we seek to use it selfishly—for
the advantage of any one nation or
any small group of nations—we
shall be equally guilty of that be-
trayal."

Fervent Interpolation

The President, speaking in the
auditorium of the War Memorial
Opera House, built in memory of
sons of the Golden Gate city who
gave their lives in the first World
War, in which he himself served,
seemed to give unconscious expres-
sion to the solemn feeling of the
delegation when, at the outset of his
speech, he interpolated the words:
'half a hope, half a prayer.'
'Oh, what a great day this can
be in history!'

Just before the plenary session
the President accompanied the
eight United States delegates to
the auditorium of the War Memorial

the Draft Ends No

WASHINGTON, Jan. 27,
1973—"With the signing of
the peace agreement in
Paris today, and after re-
ceiving a report from the
Secretary of the Army that
he foresees no need for



PATIENT PACKAGE INSERTS: A CONCEPT WHOSE TIME HAS COME?

The consumer's right to know is an irreversible and desirable trend of the Seventies. It extends, and properly, to a patient's right to know more about his or her prescription medications. One way, gaining favor, is through patient package inserts. Wisely-prepared and properly distributed when medically indicated, they could markedly improve patient knowledge and drug therapy—laudable goals by anyone's standards.

The PMA endorses these goals and will work with government, the health professions and consumers to achieve them.

The Advantages

The concept holds promise of benefits: better patient understanding of the product prescribed, better adherence to the treatment plan, and more awareness of possible side reactions.

Every doctor has had patients who fail to finish antibiotic regimens because they feel better. Some patients assume that if one tranquilizer or analgesic is good, two may be twice as good. Still others fail to report dizziness while on antihypertensive therapy—and so on.

Problems like these might arise less often if the patient received written information in addition to verbal instructions. Some studies suggest that patients are more receptive to such materials, and they more often understand the verbal instructions and follow them, when inserts are used.

The Disadvantages

There are also some potential problems. Obviously, the inserts must be clearly phrased, without extraneous or complex detail. How much information

is enough? How can it be kept current? Should all patients receive the same information? Should inserts be included with all drugs? Should only potential problems be listed or are patients better off with a "fair balance" presentation that describes usefulness as well as drawbacks?

These and similar questions require answers, since model inserts have yet to be properly developed and tested. Despite the need for these studies, the FDA is proceeding prematurely with inserts on selected products. We think the Congress is the only place where the matter can be given the proper legal status and direction, particularly since it represents a conceptual change in the legal, medical and social framework of the nation's prescription drug information system.

The Solution

The PMA believes that carefully-devised pilot studies of various kinds of inserts are needed. They should be developed and implemented with full participation by doctors, pharmacists, consumers, communications experts and the drug industry. Such studies will provide reliable pathways to follow, so that inserts will be useful aids to medical practice.

And particularly we think that you should be closely involved in this debate and in these studies and decisions. Otherwise, people with less experience and qualifications may control the purposes, content and use of a tool with considerable promise for improved patient care. It could make a difference in your practice tomorrow, and more importantly, in the health of your patients.



THE PHARMACEUTICAL MANUFACTURERS ASSOCIATION
1155 FIFTEENTH ST., N. W., WASHINGTON, D. C. 20005

Clinical experience & continuing confidence

KAON® ELIXIR was introduced in 1954, followed by KAON® TABLETS in 1963. Decades of clinical experience indicate acceptability, effectiveness, and safety in the majority of patients; should abdominal pain occur, therapy should be discontinued. Both have been taken by patient after patient, day after day, year after year, to correct potassium deficiencies. Both have consistently demonstrated their value when diet alone is inadequate for potassium replacement.

Kaon® Elixir (potassium gluconate) **Kaon® Tabs** (potassium gluconate)

BRIEF SUMMARY **Kaon Tablets/Kaon Elixir**

KAON® (potassium gluconate) TABLETS

Description: Each sugar-coated tablet supplies 5 mEq. of elemental potassium (as potassium gluconate 1.17 Gm.). Kaon Tablets are sugar coated, not enteric coated, which favors dissolution in the stomach and absorption before reaching the small intestine where the lesions with enteric potassium chloride have occurred. The sugar coating merely adds to palatability and ease of swallowing, not to delay absorption as does the enteric coating.

Indications: Oral potassium therapy for the prevention and treatment of hypokalemia which may occur secondary to diuretic or corticosteroid administration. It may be used in the

treatment of cardiac arrhythmias due to digitalis intoxication.

Contraindications: Severe renal impairment with oliguria or azotemia, untreated Addison's disease, adynamia episodica hereditaria, acute dehydration, heat cramps and hyperkalemia from any cause.

Warning: There have been several reports, published and unpublished, concerning nonspecific small-bowel lesions consisting of stenosis, with or without ulceration, associated with the administration of enteric-coated potassium tablets alone or when they are used with nonenteric-coated thiazides or certain other oral diuretics. These small-bowel lesions have caused obstruction, hemorrhage and perforation. Surgery was frequently required and deaths have occurred. Available information tends to implicate enteric-coated potassium salts, although

lesions of this type also occur spontaneously. Therefore, coated potassium-containing formulations should be administered only when indicated and should be discontinued immediately if abdominal pain, distention, nausea, vomiting, or gastrointestinal bleeding occur. Coated potassium tablets should be used only when adequate dietary supplementation is not practical.

Precautions: In response to a rise in the concentration of body potassium, renal excretion of potassium is increased. With normal kidney function it is difficult, therefore, to produce potassium intoxication by oral administration. However, potassium supplements must be administered with caution, since the amount of the deficiency or daily dosage is not accurately known. Frequent checks of the clinical status of the patient and periodic ECG and/or serum potassium levels should be made. High serum concentra-

*Time is
the test of
all things*



of potassium ion may cause death through
acid depression, arrhythmias or arrest. This
should be used with caution in the presence
of cardiac disease.

In hypokalemic states, especially in patients
on a salt-free diet, hypochloremic alkalosis is a
condition that may require chloride as well as
potassium supplementation. In these circum-
stances, Kaon (potassium gluconate) should be
supplemented with chloride. Ammonium chlo-
ride is an excellent source of chloride ion (18.7
mg. per Gram), but it should not be used in
patients with hepatic cirrhosis where ammonium
is contraindicated. Other sources for
chloride are sodium chloride and Diluted
Hydrochloric Acid, U.S.P.

It should also be kept in mind that ammonium
cation exchange resin, sometimes used to
treat hyperkalemia, should not be administered

to patients with hepatic cirrhosis.

Adverse Reactions: Nausea, vomiting, diarrhea
and abdominal discomfort have been reported.
The symptoms and signs of potassium intoxi-
cation include paresthesias of the extremities,
flaccid paralysis, listlessness, mental confusion,
weakness and heaviness of the legs, fall in
blood pressure, cardiac arrhythmias and heart
block. Hyperkalemia may exhibit the following
electrocardiographic abnormalities: disappear-
ance of the P wave, widening and slurring of
QRS complex, changes of the S-T segment, tall
peaked T waves, etc.

Overdosage: Potassium intoxication may result
from overdosage of potassium or from thera-
peutic dosage in conditions stated under
"Contraindications." Hyperkalemia, when de-
tected, must be treated immediately because
lethal levels can be reached in a few hours.

KAON® (potassium gluconate) ELIXIR

Description: Each 15 ml. (tablespoonful) sup-
plies 20 mEq. of elemental potassium (as potas-
sium gluconate, 4.68 Gm.) with saccharin and
aromatics. Alcohol 5%.

Indications: See Kaon Tablets.


Precautions: See Kaon Tablets.

In hypochloremic alkalosis, potassium
replacement with potassium chloride
(e.g., Kaochlor® 10% Liquid) may be more ad-
vantageous than with other potassium salts.

Adverse Reactions: See Kaon Tablets.

Overdosage: See Kaon Tablets.

WARREN-TEED
LABORATORIES, INC.
DIVISION OF ADRIA LABORATORIES INC.
COLUMBUS, OHIO 43215



California Medical Association
107th Annual Session
Atop Nob Hill
March 17-22, 1978
San Francisco

SPECIAL
PROGRAM

Tell it to the FDA

Saturday, March 18, 1978

2:00 to 5:00 p.m. • Masonic Temple Auditorium

The Food and Drug Administration is genuinely interested in establishing communication with practicing physicians. At the CMA Annual Session, high-level representatives of the FDA will conduct an open meeting with interested members of the CMA to foster this communication. Physicians are at the interface where the actions or interactions of the FDA have an effect upon patient care. This is an opportunity to "Tell it to the FDA."

Just Published!

THE MAKING OF A MEDICARE DOCTOR

By John M. Lanham, MD

The devastating story of how a Medicare physician must compromise his principles in order to continue to function within the bureaucratic system that demands such compromise. Reviews the implications of the current malpractice crisis and the bureaucracy in control of the free-enterprise system. **\$5.95**

At your bookstore or postpaid from:

VANTAGE PRESS, Inc.

516 W. 34 St., New York, N.Y. 10001

Tablets **Percodan®**

DESCRIPTION Each yellow, scored tablet contains 4.50 mg. oxycodone HCl (WARNING: May be habit forming), 0.38 mg. oxycodone terephthalate (WARNING: May be habit forming), 224 mg. aspirin, 160 mg. phenacetin, and 32 mg. caffeine.

INDICATIONS For the relief of moderate to moderately severe pain.

CONTRAINDICATIONS Hypersensitivity to oxycodone, aspirin, phenacetin or caffeine.

WARNINGS **Drug Dependence** Oxycodone can produce drug dependence of the morphine type and, therefore, has the potential for being abused. Psychic dependence, physical dependence and tolerance may develop upon repeated administration of PERCODAN®, and it should be prescribed and administered with the same degree of caution appropriate to the use of other oral narcotic-containing medications. Like other narcotic-containing medications, PERCODAN® is subject to the Federal Controlled Substances Act.

Usage in ambulatory patients Oxycodone may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. The patient using PERCODAN® should be cautioned accordingly.

Interaction with other central nervous system depressants Patients receiving other narcotic analgesics, general anesthetics, phenothiazines, other tranquilizers, sedative-hypnotics or other CNS depressants (including alcohol) concomitantly with PERCODAN® may exhibit an additive CNS depression. When such combined therapy is contemplated, the dose of one or both agents should be reduced.

Usage in pregnancy Safe use in pregnancy has not been established relative to possible adverse effects on fetal development. Therefore, PERCODAN® should not be used in pregnant women unless, in the judgment of the physician, the potential benefits outweigh the possible hazards.

Usage in children PERCODAN® should not be administered to children.

Salicylates should be used with caution in the presence of peptic ulcer or coagulation abnormalities.

PRECAUTIONS **Head injury and increased intracranial pressure** The respiratory depressant effects of narcotics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, other intracranial lesions or pre-existing increase in intracranial pressure. Furthermore, narcotics produce adverse reactions which may obscure the clinical course of patients with head injuries.

Acute abdominal conditions The administration of PERCODAN® or other narcotics may obscure the diagnosis or clinical course in patients with acute abdominal conditions.

Special risk patients PERCODAN® should be given with caution to certain patients such as the elderly, debilitated, and those with severe impairment of hepatic or renal function, hypothyroidism, Addison's disease, and prostatic hypertrophy or urethral stricture.

Phenacetin has been reported to damage the kidneys when taken in excessive amounts for a long time.

ADVERSE REACTIONS The most frequently observed adverse reactions include light-headedness, dizziness, sedation, nausea and vomiting. The effects seem to be more prominent in ambulatory patients in nonambulatory patients, and some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include euphoria, dysphoria, constipation and pruritus.

DOSAGE AND ADMINISTRATION Dosage should be adjusted according to the severity of the pain and response of the patient. The usual adult dose is 1 tablet every 6 hours as needed for pain.

DRUG INTERACTIONS The CNS depressant effect of PERCODAN® may be additive with that of other CNS depressants. See WARNINGS.

DEA Order Form Required.

Endo Inc.

Manati, Puerto Rico 00701
Subsidiary of Endo Laboratories, Inc.
Subsidiary of the DuPont Company



1. Determine need

What is causing pain? How is it perceived by you and your patient?

2. Prescribe a rapid-acting agent

Select a readily-absorbed oral agent that usually acts within 15 to 30 minutes.

3. Minimize potential risk

Prescribe in limited quantities for selected patients.

Schedule II classification means no refills, no telephone Rx. Patients with persistent pain must return for your evaluation of analgesic needs.

4. Provide adequate analgesia with minimum doses


Consider PERCODAN® because patients rarely ask for increased dosage. PERCODAN® relief can last up to six hours—until time for next tablet.



Effective relief of moderate to moderately severe pain

Tablets
PERCODAN®

each yellow, scored tablet contains: 4.50 mg oxycodone HCl (WARNING: may be habit forming), 0.38 mg oxycodone terephthalate (WARNING: may be habit forming), 224 mg aspirin, 160 mg phenacetin, 32 mg caffeine



PREMINE
albuterol sulfate

A second wind
for asthmatics



Information is supplied
by the manufacturer.

BRETHINE[®]
terbutaline sulfate

Geigy

It could mean a more active life for patients with reversible obstructive airways disease.

More effective

Brethine was more effective and longer acting than metaproterenol in a study of five patients with exercise-induced asthma.

Brethine has been shown to be highly effective alone and in combination with theophylline.

And Brethine has been shown to be twice as effective as ephedrine.

Long acting

Effect may last from 6 to 8 hours. One tablet at bedtime, upon arising and at midafternoon should keep patients breathing comfortably.

Minimal cardiac effect

Brethine produces proportionally greater changes in pulmonary function than in heart rate or blood pressure.

Tablets of 2.5 mg and 5 mg.

Some patients may experience mild hand tremor or "shakiness" when Brethine therapy is initiated. This may be minimized by starting patients with 2.5 mg doses.

Brethine[®], brand of terbutaline sulfate, Tablets 5 mg., Tablets 2.5 mg. Before prescribing or administering, please consult complete product information, a summary of which follows:

Tablets contain 5 mg. (equivalent to 4.1 mg. of free base) or 2.5 mg. (equivalent to 2.05 mg. of free base) of Brethine, brand of terbutaline sulfate.

Indications: As a bronchodilator for bronchial asthma and for reversible bronchospasm which may occur in association with bronchitis and emphysema.

Contraindications: Known hypersensitivity to sympathomimetic amines.

Warnings: *Usage in Pregnancy:* The safety of the use of Brethine, brand of terbutaline sulfate, in human pregnancy has not been established. The use of the drug in pregnancy, lactation, or women of childbearing potential requires that the expected therapeutic benefit of the drug be weighed against its possible hazards to the mother or child.

Usage in Pediatrics: Brethine, brand of terbutaline sulfate, tablets are not presently recommended for children below the age of twelve years due to insufficient clinical data in this pediatric group.

Precautions: Brethine, brand of terbutaline sulfate, should be used with caution in patients with diabetes, hypertension, and hyperthyroidism. As with other sympathomimetic bronchodilator agents, Brethine, brand of terbutaline sulfate, should be administered cautiously to cardiac patients, especially those with associated arrhythmias. Although the concomitant use of Brethine, brand of terbutaline sulfate, with other sympathomimetic agents is not recommended, the use of an aerosol bronchodilator of the adrenergic stimulant type for the relief of an acute bronchospasm is not precluded in patients receiving chronic oral Brethine, brand of terbutaline sulfate, therapy.

Adverse Reactions: Commonly observed side effects include nervousness and tremor. Other reported reactions include headache, increased heart rate, palpitations, drowsiness, nausea, vomiting, and sweating. These reactions are generally transient in nature, usually do not require treatment, and appear to diminish in frequency with continued therapy. In general, all the side effects observed are characteristic of those commonly seen with sympathomimetic amines.

How Supplied: Round, scored, white tablets of 5 mg. in bottles of 100 and 1,000 and Unit Dose Packages of 100; oval, scored, white tablets of 2.5 mg. in bottles of 100. (B) 98-146-060-E (Rev. 4/76) 667004 C76-12

GEIGY Pharmaceuticals
Division of CIBA-GEIGY Corporation
Ardsley, New York 10502

**Brethine: alone or with theophylline, a second wind
for asthmatics.**

Personally, Steve could never speak up for milk.

In TV commercials, you'll often see people speaking up for milk. Steve Allen and his wife Jayne Meadows are also seen on TV telling viewers about Mocha Mix...100% milk free!


Matter of fact, this non-dairy creamer would have been found in their home way before they ever filmed the Mocha Mix series of commercials. Steve is allergic to milk products. Yet, he likes the splash of luxury and fresh taste in his coffee, on cereal, fruit and dessert, even in cooking. For the Allen household, Mocha Mix is the perfect answer.

It could well be the answer for many people who



rely on you for their dietary needs, those with an allergy similar to Steve's, or a lactose intolerance. And, for many who must maintain a fat-restricted diet you should know that Mocha Mix is one non-dairy creamer that contains no coconut oil, giving it

the highest ratio of unsaturated to saturated fat of any creamer. It easily exceeds the accepted standard 2:1 ratio. It's 100% cholesterol free, too. No wonder Mocha Mix is not just another non-dairy creamer, but the one you can rely on most for your patients who can't speak for milk, either.



NUTRITION INFORMATION

Portion Size 1 Fluid Ounce (2 Tbs.)

Servings Per Container	16
Calories	40
Protein	0 grams
Carbohydrate	3 grams
Fat	3 grams
Percent of Calories from Fat	73%
Polyunsaturated Fat	1 gram
Saturated Fat	0 grams
Cholesterol (A)	0 mg (0 mg/100g)

In addition to the pint and quart size found in the dairy case of most grocery stores, Mocha Mix is available in 4 ounce and 1/2 oz. portion packs for hospitals and institutions.

Interested? Send us a note and we will send you a supply of coupons your patients can redeem at their grocers. Hospital service may also be supplied upon request. Mail to: Mocha Mix Dept. Presto Food Products, Inc. P.O. Box No. 21908, Los Angeles, Calif. 90021

Percentage of U.S. Recommended Daily Allowance (U.S. RDA)*

* Contains less than 2% of the U.S. RDA of Protein, Vitamin A, Vitamin C, Thiamine, Riboflavin, Niacin, Calcium, Iron.

Information on fat and cholesterol content is provided for individuals who, on the advice of a physician, are modifying their total dietary intake of fat and cholesterol.

mocha mix®

... the non-dairy creamer that's lowest in saturated fat.

TRIAMTERENE CONSERVES POTASSIUM WHILE HYDROCHLOROTHIAZIDE LOWERS BLOOD PRESSURE **DYAZIDE®**

Each capsule contains 50 mg. of Dyrenium® (triamterene, SK&F Co.) and 25 mg. of hydrochlorothiazide.

MAKES SENSE

Before prescribing, see complete prescribing information in SK&F Co. literature or PDR. A brief summary follows:

* Warning

This drug is not indicated for initial therapy of edema or hypertension. Edema or hypertension requires therapy titrated to the individual. If this combination represents the dosage so determined, its use may be more convenient in patient management. Treatment of hypertension and edema is not static, but must be reevaluated as conditions in each patient warrant.

* **Indications:** When the combination represents the dosage determined by titration: Adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, the nephrotic syndrome. Corticosteroid and estrogen-induced edema, idiopathic edema; hypertension, when the potassium sparing action of triamterene is warranted. (See Box Warning.) Routine use of diuretics in healthy pregnant women is inappropriate; they are indicated in pregnancy only when edema is due to pathological causes.

Contraindications: Further use in anuria, progressive renal or hepatic dysfunction, hyperkalemia. Pre-existing elevated serum potassium. Hypersensitivity to either component or other sulfonamide-derived drugs.

Warnings: Do not use potassium supplements, dietary or otherwise, unless hypokalemia develops or dietary intake of potassium is markedly impaired. If supplementary potassium is needed, potassium tablets should not be used. Hyperkalemia can occur has been associated with cardiac irregularities. It is more likely in the severely ill, with urine volume less than one liter/day, the elderly and diabetics with suspected or confirmed renal insufficiency. Periodically, serum K^+ levels should be determined. If hyperkalemia develops, substitute a thiazide alone, restrict K^+ intake. Associated widened QRS complex or arrhythmia requires prompt additional therapy. Thiazides cross the placental barrier and appear in cord blood. Use in pregnancy requires weighing anticipated benefits against possible hazards, including fetal or neonatal jaundice, thrombocytopenia, other adverse reactions seen in adults. Thiazides appear and triamterene may appear in breast milk. If their use is essential, the patient should stop nursing. Adequate information on use in children is not available.

Precautions: Do periodic serum electrolyte determinations (particularly important in patients vomiting excessively or receiving parenteral fluids).

Periodic BUN and serum creatinine determinations should be made, especially in the elderly, diabetics or those with suspected or confirmed renal insufficiency. Watch for signs of impending coma in severe liver disease. If spironolactone is used concomitantly, determine serum K^+ frequently; both can cause K^+ retention and elevated serum K^+ . Two deaths have been reported with such concomitant therapy (in one, recommended dosage was exceeded, in the other serum electrolytes were not properly monitored). Observe regularly for possible blood dyscrasias, liver damage, other idiosyncratic reactions. Blood dyscrasias have been reported in patients receiving triamterene, and leukopenia, thrombocytopenia, agranulocytosis, and aplastic anemia have been reported with thiazides. Triamterene is a weak folic acid antagonist. Do periodic blood studies in cirrhotics with splenomegaly. Antihypertensive effect may be enhanced in post-sympathectomy patients. Use cautiously in surgical patients. The following may occur: transient elevated BUN or creatinine or both, hyperglycemia and glycosuria (diabetic insulin requirements may be altered), hyperuricemia and gout, digitalis intoxication (in hypokalemia), decreasing alkali reserve with possible metabolic acidosis.

'Dyazide' interferes with fluorescent measurement of quinidine.

Adverse Reactions:

Muscle cramps, weakness, dizziness, headache, dry mouth; anaphylaxis, rash, urticaria, photosensitivity, purpura, other dermatological conditions;


nausea and vomiting, diarrhea, constipation, other gastrointestinal disturbances. Necrotizing vasculitis, paresthesias, icterus, pancreatitis, xanthopsia and, rarely, allergic pneumonitis have occurred with thiazides alone.


Supplied: Bottles of 100 and 1000 capsules; Single Unit Packages of 100 (intended for institutional use only).

**FOR LONG-TERM CONTROL
OF HYPERTENSION.*
SERUM K^+ AND BUN SHOULD
BE CHECKED PERIODICALLY.
(SEE WARNINGS SECTION.)**

SK&F CO., Carolina, P.R. 00630

SK&F CO.
a SmithKline company





Simple
drops of water
help make
COLACE the
most widely used
stool softener.

COLACE®

docusate sodium sulfosuccinate

COLACE works by stool-softening action, free from laxative stimulation. Simply by letting natural intestinal action do most of the work, COLACE helps to eliminate the hard, dry stools common to constipation.

COLACE works to prevent pain and straining at stool with minimum chance of griping or cramps, particularly in patients with delicate anorectal disorders. COLACE is safe and non-habit forming in short- or long-term therapy. COLACE—the simple water way to ease constipation from infancy to old age. Capsules, syrup or liquid.

Mead Johnson PHARMACEUTICAL DIVISION

COMPATIBILITY



Eastern Tiger Swallowtail Butterfly
(*Papilio glaucus*)

Does it influence your choice of peripheral/cerebral vasodilator*?

Vasodilan—compatible with coexisting diseases (e.g., glaucoma, diabetes)

Vasodilan has not been reported to affect the course of coexisting disease; it has not been reported to affect blood sugar levels or to raise intraocular pressure.

Vasodilan—compatible with concomitant therapy

Vasodilan has not been reported to affect the treatment of coexisting disease; it is compatible with such drugs as hypoglycemics and miotics.

Vasodilan—compatible with your total regimen for peripheral vascular insufficiency

Vasodilan can be a valuable adjunct in planning a total therapeutic program for peripheral vascular insufficiency.

Indications: Based on a review of this drug by the National Academy of Sciences-National Research Council and/or other information, the FDA has approved the indications as follows:

Indications Effective:

For the relief of symptoms associated with cerebral vascular insufficiency, peripheral vascular disease of arteriosclerosis obliterans, thromboangiitis obliterans (Buerger's Disease) and Raynaud's disease. Further classification of the less-than-effective indications requires further investigation.

Dosage: Vasodilan tablets, isoxsuprine HCl, 10 mg. and 20 mg. Vasodilan injection, isoxsuprine HCl, 5 mg., per ml.

Mode and Administration: Oral: 10 to 20 mg., three or four times daily. Intramuscular: 5 to 10 mg. (1 or 2 ml.) two or three times daily. Intramuscular administration may be used initially in severe or acute conditions.

Contraindications and Cautions: There are no known contraindications to oral administration in recommended doses. Should not be given immediately after a meal or in the presence of arterial bleeding.

Parenteral administration is not recommended in the presence of hypotension or tachycardia.

Intravenous administration should not be given because of increased likelihood of side effects.


Adverse Reactions: On rare occasions oral administration of the drug has been associated in time with the occurrence of hypotension, tachycardia, nausea, vomiting, dizziness, abdominal distress, and severe rash. If rash appears the drug should be discontinued.

Although available evidence suggests a temporal association of these reactions with isoxsuprine, a causal relationship can be neither confirmed nor refuted.

Administration of single dose of 10 mg. intramuscularly may result in hypotension and tachycardia. These symptoms are more pronounced in higher doses. For these reasons single intramuscular doses exceeding 10 mg. are not recommended. Repeated administration of 5 to 10 mg. intramuscularly at suitable intervals may be employed.

Supplied: Tablets, 10 mg., bottles of 100, 1000, 5000 and Unit Dose; Tablets, 20 mg., bottles of 100, 500, 1000, 5000 and Unit Dose; Injection, 10 mg. per 2 ml. ampul, box of six 2 ml. ampuls.

U.S. Pat. No. 3,056,836




VASODILAN[®] 20-mg tablets

(ISOXSUPRINE HCl)

20 mg q.i.d. recommended dosage

Mead Johnson
PHARMACEUTICAL DIVISION

© 1978 MEAD JOHNSON & COMPANY • EVANSVILLE, INDIANA 47721 U.S.A. MJL 7-4237R



**This asthmatic
isn't worried about his next breath...**

**he's active
he's effectively
maintained on**

QUIBRON

Each capsule or tablespoon (15 ml) elixir contains theophylline (anhydrous) 150 mg and glyceryl guaiacolate (guaifenesin) 90 mg. Elixir: alcohol 15%

- high theophylline for effective around-the-clock therapy
- 100% free theophylline
- individualized theophylline dosage schedule

Indications: For the symptomatic treatment of bronchospastic conditions such as bronchial asthma, asthmatic bronchitis, chronic bronchitis, and pulmonary emphysema.

Dosage: Initial: Adults: 1-2 capsules or 1-2 tablespoons elixir every 6-8 hours, children 8-12: 1 tablespoonful or one capsule every 6-8 hours and children under 8: 3 to 5 mg theophylline/kg body weight every 6-8 hours. Theophylline dosage may be cautiously increased to 2000 mg/24 hr in adults or 7 mg/kg in children; monitoring of serum theophylline levels at higher dosages is recommended.

Precautions: Do not administer more frequently than every 6 hours, or within 12 hours after rectal dose of any preparation containing theophylline or aminophylline. Do not give other xanthine derivatives concurrently. Use in case of pregnancy only when clearly needed.

Adverse Reactions: Theophylline may exert some stimulating effect on the central nervous system. Its administration may cause local irritation of the gastric mucosa, with possible gastric discomfort, nausea and vomiting. The frequency of adverse reactions is related to the serum theophylline level and are not usually a problem at serum theophylline levels below 20 µg/ml.

How Supplied: Capsules in bottles of 100 and 1000 and unit-dose packs of 100; Elixir in bottles of 1 pint and 1 gallon.

Mead Johnson

PHARMACEUTICAL DIVISION

© 1978 Mead Johnson & Company • Evansville, Indiana 47721 U.S.A. MJL 6-78

Kefzol® I.M./I.V. cefazolin sodium

Ampoules, equivalent to 500 mg., 1 Gm.,
and 10 Gm. of cefazolin



700773

Additional information available
to the profession on request.
Eli Lilly and Company
Indianapolis, Indiana 46206

Classified Advertisements

The rate for each insertion is **\$2.50 per line** (average six words per line) with **five line minimum**.

Box number charge: **\$1.50 each month**.

Classified display rates \$25.00 per inch.

Copy for classified advertisements should be received not later than the **fifth of the month preceding issue**. • Classified advertisers using Box Numbers forbid the disclosure of their identity. Your inquiries in writing will be forwarded to Box Number advertisers. The right is reserved to reject or modify all classified advertising copy in conformity with the rules of the Advertising Committee.

CLASSIFIED ADVERTISEMENTS ARE PAYABLE IN ADVANCE

PHYSICIANS WANTED

CARDIOLOGIST, certified or recently eligible, wanted for expanding 47 member multispecialty group with 16 internists, most having subspecialty certifications, in rapidly growing 95,000 city, 90 miles east of San Francisco. Invasive and non-invasive cardiology labs, open heart surgery in local hospitals. Competitive compensation with excellent fringe benefits. California license required. Reply with curriculum vitae to Darrell K. Oberg, MD, The Gould Medical Group, Inc., 600 Coffee Road, Modesto, CA 95355. (209) 524-1211.

ANESTHESIOLOGIST—California, Board Certified, for Chief, Anesthesia Section, needed at VA Hospital, Livermore, CA. Ideal living, good climate and clean air. Malpractice insurance not needed. Position is available at present time. Salary dependent on experience and qualifications. Contact: Byron V. Whitney, MD, FACS, Chief, Surgical Service, Veterans Administration Hospital, Livermore, CA 94550. Tel: (415) 447-2560, ext. 213.

MEDICAL DIRECTOR—Fresno County Department of Health, Outpatient Family Planning Facility desires a full-time Medical Director. Forty hour week. No abortions. Contract \$48,000 depending on qualifications and experience. Vacation, paid malpractice, and educational leave provided. Staff privileges at teaching hospital available to appropriate candidate. Board certified OB-Gyn (or eligible). Fresno is a delightful city of 186,000 located three hours by car from San Francisco, Los Angeles, Yosemite, and Pacific Coast. Write Ms. Zoe Ann Conley, P.O. Box 11867, Fresno, CA 93775, or call (209) 488-3004 collect.

OVERSEAS OPPORTUNITY PHYSICIAN FOR MID-EAST ASSIGNMENT TO PROVIDE HEALTH CARE TO EXPATRIATES

Family practitioner or emergency room physician with 4 years or more experience, board certification desirable. Excellent salary and fringe benefits; liberal R&R program with opportunity to travel.

Single status assignment, one-year contract. Management and administrative skills along with competent medical expertise desirable. Responsible position; must be capable of working independently and supervise allied health professionals.

INTERNATIONAL MEDICAL SERVICES

630 Grovenor Plaza
150 South Los Robles Avenue
Pasadena, California 91101
(213) 577-0940

INTERNIST-GENERAL PRACTITIONER sought by Medical Service, Brentwood Veterans Administration Hospital. Serves as Consultant to the hospital and as one of several staff physicians on twenty-five bed inpatient/outpatient medical service. Acute general, respiratory, gastrointestinal, hematologic, and neurologic problems are treated. All Surgical, Radiological, Laboratory, and sub-specialty Medical Consultation services are available. Close affiliation with UCLA; qualified internists can seek appointment to the Clinical Faculty of UCLA department of Medicine. Team approach provides support of other internists, excellent nursing staff, social work. Write Milton Greenblatt, MD, Chief of Staff, VA Hospital Brentwood, Wilshire and Sawtelle Blvds., Los Angeles, CA 90073—Phone (213) 478-3711, Ext. 2641. Equal Opportunity Employer.

FP's for fresh approach to Primary Care. Group practice with independent freedom. All administration eliminated. No financial risks. Hospital-based or associate practices in several pleasant N. California communities. FFS with \$40,000 annual guarantee. Minimal night-call. E.R. back-up available. **CONTACT: EMS, 2310 Mason St., San Francisco, CA 94133. (415) 956-5900.**

TEACHING POSITION FOR EXPERIENCED FAMILY PHYSICIAN—Position available as director of the Williamsburg Family Practice Office, a teaching practice, affiliated with the Department of Family Practice, University of Iowa. American Board of Family Practice certification required. Prosperous farming community, practice serves 6,000 people. New office building. 25 miles from University Medical Center. Faculty appointment with salary and University fringe benefits, including malpractice insurance. University community offers excellent cultural and educational opportunities. Night and weekend coverage will be shared by residents. Direct inquiries to: Robert E. Rakel, MD, Professor and Head, Department of Family Practice, University of Iowa College of Medicine, Iowa City, Iowa 52242.

GERIATRICIAN—Physician experienced in geriatrics and geriatric programs; to design and implement inpatient and outpatient geriatrics program; participate in approved internal medicine and surgery residency training programs; provide direct patient care. Send résumé to Russell D. Tyler, MD, Medical Director, Health Care Services, Santa Barbara General Hospital, P.O. Box 3650, Santa Barbara, CA 93105.

INTERNIST NEEDED for multispecialty group. Excellent working conditions, fringe benefits, salary and bonus dependent on individual effort. Excellent medical community. Maintenance of individuality within the corporate structure stressed. Interested parties contact F. E. Roque, MD, Fresno Medical Group, Inc., 1300 N. Fresno St., Fresno, CA 93703. Phone (209) 486-3610.

CALIFORNIA, SAN JOSE: Experienced, career-oriented Emergency Physicians wanted to join well-established group practicing at a university affiliated hospital and a large community hospital. Beautiful San Francisco Bay area location. Fee-for-service compensation. ACEP preferred. Contact: James B. Lane, MD at (408) 293-8881 or write: 1530 The Alameda, No. 28, San Jose, CA 95126.

BOARD CERTIFIED OR ELIGIBLE—General Internist, Oncologist needed for 45-man multispecialty group. Adjacent 400-bed community hospital. Excellent location, San Francisco Bay Area. University teaching affiliations available. Contact Administration, The San Jose Medical Clinic, 45 So. 17th Street, San Jose, CA 95112. Phone (408) 998-5551.

SHASTA COUNTY—REDDING. Shasta General Hospital (fully accredited) has vacancy for board certified or eligible OB/GYN specialist. Active outpatient clinic and inpatient unit. 12 full-time physician staff includes 2 OB/GYN. \$43,000 yr. (\$44,500 if board certified). Excellent benefits include prepaid malpractice insurance. Contact Eugene Earl, MD, Medical Director, Shasta General Hospital, 2630 Hospital Lane, Redding, CA 96001. Phone (916) 241-3232.

(Continued on Page 24)



ASSISTANT DIRECTOR MEDICAL AFFAIRS SAUDI ARABIA

A unique overseas challenge offering travel, an excellent salary and personal enrichment.

Whittaker Corporation, a leader of international health care programs, is currently seeking an individual with solid experience in U.S. hospitals and medical administration with the ability to establish standards to measure and audit the quality of medical and hospital care. Will be responsible for evaluating and making recommendations on medical and administrative matters, in addition to participation in medical care activities at various hospital sites.

Ideal candidate should be a graduate of an accredited medical school with a minimum of three to five years as department head with increasing responsibilities and a proven track record of administrative abilities including excellent communication skills.

Our overseas compensation includes a competitive salary and benefits package with furnished housing, substantial vacation, travel opportunities and other attractive fringe benefits.

For further information and confidential consideration, please forward your curriculum vitae to:

LARRY M. ROSS
Manager, International Recruitment

Whittaker

LIFE SCIENCES GROUP

10880 Wilshire Boulevard, Suite 604,
Los Angeles, California 90024

An Equal Opportunity Employer M/F

ANESTHESIOLOGISTS

There is a better way for solo or group practice receivables management.

Anesthesiologist Services, Incorporated
"The Cash Flow Specialists"
Call (800) 652-1450 or (415) 441-8186 or write
P.O. Box 29292, San Francisco, CA 94128

PHYSICIAN NEEDED

A relaxing atmosphere but rewarding challenge for 2 G.P.'s preferably surgery and OB experience, is waiting for you. All outdoor activities and sports are available for all seasons, cultural arts are within 40 miles. A medical school and continuing education is closeby and available. The hospital has 31 beds with 3 G.P.'s on staff. A doctor's clinic is available immediately adjacent to the hospital plus other locations.

Please send CV or call collect:

(801) 623-1242
David Peterson, Administrator
Juab County Hospital
Nephi, Utah 84648

“Revelation of a second death, and new animal studies, indicate Laetrile can cause fatal poisoning...Georgia researchers will test laser beams for arthritis treatment...In Alabama, a vaccine for protection against dental caries is being studied...New York and Swedish clinicians unveil two noninvasive screening methods for detection of breast cancer...”

Important things are happening that you should know about right away. You'll find them on your desk every two weeks in Medical World News, the newsmagazine of medicine.

Read this one first.

medical world news

A McGraw-Hill publication



PHYSICIANS WANTED

STUDENT HEALTH-MEDICAL OFFICER II, ASSISTANT CLINICAL DIRECTOR. Full time, 12 months position available, July 1, 1978. Participate in health care delivery, help supervise clinical personnel and also assist the Director in some administrative areas. Family Practice, Internal Medicine, or Pediatric background preferred and desirable. Interest in student-age health care and wellness essential. We have an exceptional staff and facilities for student health care at California State University, Fullerton! Twenty-two thousand students and 2,500 faculty and staff; near to beaches, mountains and desert. Range of salary—\$34,728-\$42,024, annually. Send résumé to Robert J. McFerran, MD, Medical Director, Student Health Service, California State University, Fullerton, 800 N. State College Blvd., Fullerton, California 92634. Applications close March 15, 1978. Equal opportunity, affirmative action employer.

PHYSICIANS WANTED—Beverly Hills, Calif.—OPHTHALMOLOGIST interested in having a well-trained man to share space in office, who is interested, ultimately, toward a partnership or eventually succeeding to this practice on mutually acceptable terms. Contact Box No. 5964, Western Journal of Medicine, 731 Market Street, San Francisco, CA 94103.

OBSTETRICIAN, SAN DIEGO—The Maternal Health Care Access Team Program has an opening for a Board Certified Obstetrician (eligible for licensure in the State of California). This is an excellent opportunity to join an innovative maternity care program that serves patients in both a Major Medical Center and a Community Clinic setting. Salary is negotiable; medical liability coverage is provided. Flexible schedule. Area located close to beaches, mountains, and many recreational facilities. Direct curriculum vitae to Patricia L. Mahorney, Project Director, Mercy Hospital and Medical Center, 4077 Fifth Avenue, San Diego, CA 92103, or phone (714) 294-8565.

THE HAYWARD, CALIFORNIA, KAISER-PERMANENTE MEDICAL CENTER RADIOLOGY DEPARTMENT will be in need of an experienced Board Certified General Diagnostic Radiologist on or about July 1, 1978. Inquire from Chief of Radiology, Kaiser Permanente Medical Center, 27400 Hesperian Blvd., Hayward, CA 94545.

RESIDENT-SURGERY—Second post graduate year in surgery available July 1978. Fully approved 5-year program in General Surgery. No pyramid through fifth year. University affiliated in large western metropolitan area. For info, contact: Chief of Surgery, Dr. Buford Burch, Highland Hospital, 1411 E. 31st St., Oakland, CA 94602.

CHRIST HOSPITAL



OAK LAWN, ILLINOIS

DIRECTOR FAMILY PRACTICE PROGRAM FAMILY PRACTICE CENTER

Christ Hospital, an 830-bed university-affiliated teaching hospital located in southwestern suburban Chicago, seeks a full-time Director of the Family Practice Program for Christ Hospital, Oak Lawn, Illinois.

Position offers a unique opportunity for leadership in directing an expanded family practice residency program. Position includes participation in an innovative health care delivery system. Offers a challenging environment for developing teaching programs at all levels of service and medical education in a large well-organized Family Practice Center (14,000 square feet).

Qualified candidate will be board-certified in family medicine with proven experience in medical education, private patient care and administration.

Qualified candidate will be eligible for faculty appointment with affiliated Rush University School of Medicine, Chicago, Illinois. Candidate will also be part of the Evangelical Hospital Association multi-hospital system which includes Good Samaritan Hospital, Downer's Grove, and Good Shepherd Hospital, Barrington.

Salary negotiable. Complete benefit package. Interested applicants should submit curriculum vitae to:

FAMILY PRACTICE SEARCH COMMITTEE

**Christ Hospital
4440 W. 95th Street
Oak Lawn, Illinois 60453**

For more information, call

**Otto M. Janke, Vice President/Administrator
(312) 425-8000**

Hesitant About Relocating?

No need to be. American Medico's Professional Relations Department has provided assistance to hundreds of physicians who've relocated to one of the more than 50 communities we serve. If you are considering relocating to a community where there is a more challenging opportunity, or if you simply desire a new lifestyle in a different climate and geographic surrounding, get in touch with us. There is no obligation, no cost. We will help you as we have so many others. We know the geography, the economics, the demographics, and we know the physician's needs. Send a current CV or telephone the Director:

**Professional Relations Department
American Medico Hospital Services
2550 Walnut Hill Lane
Dallas, Texas 75229
(214) 358-5611**

PHYSICIAN SPECIALIST

Harbor General Hospital, Office of the Medical Director, is accepting applications for the position of Physician Specialist to supervise the Quality Assurance Program. Applicants must be Board Certified or Eligible in Internal Medicine, Pediatrics, or Family Practice.

Contact: Marvin Gasster, MD

**HARBOR GENERAL HOSPITAL
1000 W. Carson Street
Torrance, CA 90509**

PHYSICIANS

PICK A PRACTICE

If you're not where you want to be, then let us help you pick a practice.

Hospital Corporation of America has undertaken a free, no obligation community service project to help match fine talents and beautiful places. We have opportunities from Miami to San Francisco with many stops in between. Each community is equipped with modern JCAH accredited hospitals, and a recognized need for additional physicians.

If you're tired of where you are, let us help place you where you want to be. Please write us and include your curriculum vitae, starting availability date, geographical preference, type of practice, and size of community desired.

Address inquiries to:

**Kenneth W. Oliver
Director, Professional Relations**



**HOSPITAL
CORPORATION
OF AMERICA**

**One Park Plaza
Nashville, Tennessee 37203**

PHYSICIANS WANTED

MAJOR MIDWESTERN TEACHING AND RESEARCH CENTER has a career opportunity for a plastic surgeon. This appointment affords the benefits of practicing as a member of a "closed" group practice, with representatives from all medical specialties. Facilities include a 1,100 bed teaching hospital with a large outpatient clinic and three suburban satellite clinics. Major medical school affiliations afford additional teaching opportunities. Individual based salary, commensurate with experience and comprehensive benefits are offered. These include: malpractice insurance, retirement plan, BC/BS Major Medical, life, accident, travel and dental insurance coverage, as well as a lease car plan and many extras. Interested parties should send their curriculum vitae in strict confidence to Box 5966, The Western Journal of Medicine, 731 Market Street, San Francisco, CA 94103.

CHALLENGING OPPORTUNITY AVAILABLE as chairman of the newly established Department of Cellular and Molecular Biology at Henry Ford Hospital. This department's main objective is the pursuit of new knowledge and understanding of those areas of science which traditionally have been aligned to the clinical practice of medicine. The main commitment will be to pursue research in biomedicine. Responsibilities of office: (1) Direct the Department and the allocation of resources (budget, space, personnel); (2) Act as liaison to clinic departments, and serve on the Council of the Henry Ford Hospital; (3) Recruit research and teaching staff, and establish research objectives; (4) Assist in grantsmanship; (5) Facilitate teaching of allied health and medical education in coordination with the Director of Medical Education. The successful candidate for the chairmanship of this department must have a Ph.D. and/or M.D. degrees, and established a research program and a record of research productivity in the science, basic to medical care. Henry Ford Hospital, 2799 West Grand Boulevard, Detroit, Michigan 48202, Attn: Mr. Jerry Dutkewych.

LICENSED PHYSICIAN wanted to work with two other physicians in family practice and internal medicine setting in growth area with good future opportunities as well. May require some travel. Good income and ample free time. Call (916) 55-8480 or write Box 5968, The Western Journal of Medicine, 731 Market St., San Francisco, CA 94103.

SITUATIONS WANTED

OTOLOGYNGOLOGIST—32, Military completed; California university trained in all aspects of Head and Neck, Facial Plastic and Reconstructive Surgery and Otolaryngology including Maxillofacial trauma; Seeks position in Western States, Pacific Coastal preferred. Available July 1, 1978. Write Box 5965, The Western Journal of Medicine, 731 Market St., San Francisco, CA 94103.

PHYSICIAN'S ASSISTANT SEEKS EMPLOYMENT—Begin Oct. 16, 1978. Training at USC School of Emergency Medicine. Licensed as R.N., Med. Tech. (ARRT-CRT). Write Mr. Glenn Phillips, 1008 So. Valencia, Alhambra, CA 91801.

PHYSICIAN'S ASSISTANT with experience in emergency medicine and surgery desiring position in Northern California. Résumé on request. Trained at USC. Write Box 5970, The Western Journal of Medicine, 731 Market St., San Francisco, CA 94103.

PHYSICIAN UNDERWRITER seeks full-time or part-time position in San Francisco. Available after 7/78. Write Box 5967, The Western Journal of Medicine, 731 Market St., San Francisco, CA 94103.

INTERNIST-PULMONARY, board eligible, university trained. B. J. Medical Poona (India). Flex NY, NJ. Seeks solo, group, hospital-based practice or associate with retiring physician in California or Arizona or other western states. Available July '78. Ruston Damania, 108 Oak St., Ridgewood, NJ 07450. (201) 444-0487.

CERTIFIED PHYSICIAN ASSISTANTS FOR MIDDLE-EAST ASSIGNMENT TO PROVIDE CARE TO EXPATRIATES

Knowledgeable P.A. with laboratory and x-ray capabilities in addition to clinical expertise. Single status assignment. (Must be bilingual in French-English.) One year contract. Excellent salary and fringe benefits; liberal R&R program with opportunity to travel. Responsible position for person who can work independently.

INTERNATIONAL MEDICAL SERVICES

630 Grosvenor Plaza
150 South Los Robles Avenue
Pasadena, California 91101
(213) 577-0940

ANESTHESIA SERVICE GROUP

Complete coverage for hospital, clinic or office on "AS NEEDED" basis in Bay Area, So. Calif. and San Diego. Call or write to: P.O. Box 6275, Oakland CA 94603 (415) 763-0336; P.O. Box 1102, S. Monica, CA 90406 (213) 393-4267; P.O. Box 2828, S. Diego, CA 92112 (714) 231-0123.

PRACTICE FOR SALE

PRIVATE PSYCHIATRIC PRACTICE FOR SALE: Fullerton, individual and family therapy with potential for group psychotherapy. Practice established over twenty years in Fullerton with easy access to metropolitan Los Angeles, Disneyland and other beach, mountain and desert resort areas. Located within a few minutes of three general hospitals and at least ten to fifteen minutes away from in-patient psychiatric facilities. Write to 4792 Logana Plaza, Yorba Linda, CA 92686, or call (714) 970-1477.

FULLY EQUIPPED EAR, NOSE AND THROAT OFFICE in Medical Arts Building adjacent to Doctors' Hospital, San Leandro. Write Box 5969, The Western Journal of Medicine, 731 Market St., San Francisco, CA 94103.

TRANSLATOR

PHYSICIAN, European background, will translate French, Spanish, Italian into English accurately, rapidly, reasonably: M.C., 2014 N. Hoover St., Los Angeles, CA 90027. (213) 663-2614.

EQUIPMENT FOR SALE

RADIUM, nine 10 mg. platinum tubes, applicators and accessories. \$3,000 (List price \$6,000). R. N. Todd, MD, 2426 Buhne, Eureka, CA 95501, phone (707) 442-3704.

OFFICE TO SHARE

BERKELEY OFFICE, 10/20 hours weekly. Near hospital. Ideal for surgeon or specialties. (415) 848-8451.

OFFICE SPACE AVAILABLE

GROWING AREA OF DUNSMUIR, CALIFORNIA, NEEDS ANOTHER DOCTOR. Space available in Travelers Hotel. Pharmacy across street; hospital eight miles away in Mt. Shasta. Nice scenic area with clean air, water. Phone or write: Ted Santos, 425 Bush St., Suite 200, San Francisco, CA 94108. Phone (415) 391-7177.

ZANE GREY NEW ZEALAND FISHING TOUR

Historically authentic itinerary, limited to 18 persons. Combination big game and fresh water fishing plus preparatory instructional seminar presented by experts. First departure April 1978.

For Details:

TRANS-PACIFIC
SPECIAL INTEREST TOURS
P.O. BOX 40165
SANTA BARBARA, CA 93103
TEL. (805) 682-6191

VACATION RENTALS

HAWAII RENTAL—HAWAIIAN (Hanalei, Kauai) VACATION beach home for only \$250 per week, monthly rate \$950. Old Hawaiian atmosphere, away from crowded beaches with lots of open space. Excellent skin diving, swimming and beaches. Sleeps 11. For details, pictures and information, write: James R. Christiansen, 1216 State Street, Santa Barbara, CA 93101, tel. (805) 962-8141.

OFFICES TO RENT OR LEASE

SAN JOSE—New medical suites, custom designed, central air and vacuum from 1,000 sq. ft. Located in fast growing East San Jose. 500+ dwellings currently under construction. Contact Cecil Compton, agent. (408) 287-7900.

MARIN COUNTY, CALIF.—Established medical office available in downtown San Anselmo, 777 square feet. Phone (415) 453-7501.

DOWNNEY, CALIFORNIA—Completely equipped 1,100-sq.ft. medical suite. Well located, ample free parking, across from Downey Community Hospital. Phone (213) 861-0396.

MAMMOTH LAKES, CALIFORNIA medical suite. High Sierra summer and ski resort. Partially furnished, including x-ray, diathermy, gyn. room and minor surgery. Pharmacy adjacent, approximately 1,650 sq. ft. Local hospital near completion. Reduced rent first 6 months. Bruce Lance, (714) 983-9574, 1970 Morgan Ave., Claremont, CA 91711.

CLASSIFIED

INFORMATION

(415) 777-2000

EX 220


Medicine's men on the Hill

Just who are they? They're the AMA's permanent representatives to the Congress of the United States from the AMA's Washington office.

In every Congress about 10% of all legislation introduced is health related — some 2,500 bills and the number is increasing every year. The AMA lobbyists serve as the profession's eyes, ears, and voice on the Hill. Keeping in day-to-day contact with the members of Congress and their staffs. Explaining and promoting the profession's views. Reporting on legislation. And providing legislators with resource material and information on medical and health subjects.

They're on the Hill to protect your interests, lobbying to retain the basic freedoms of medical practice in any government health program that might be enacted. Equally important, they lobby to insure the passage of constructive and workable health legislation for the public.

Sure, the AMA lobbies. It lobbies for the rights and interests of our profession and for quality medical care for every American. With your support, the AMA can be even more effective.



Join us.
We can do much more together.

Dept. of Membership Development
American Medical Association
535 N. Dearborn St./Chicago, IL 60610

Please send me more information on the AMA
and AMA membership.

Name _____

Address _____

City/State/Zip _____

If you want your patients to buy the margarine highest in beneficial polyunsaturates, recommend Saffola®.

Saffola margarine is made with liquid safflower oil. So it's higher in polyunsaturates than even corn oil margarine.

And Saffola margarine contains no cholesterol. But it tastes as delicate and light as other spreads.

Along with Saffola mayonnaise and Saffola safflower oil,


Saffola margarine can help your patients reduce serum cholesterol as part of a fat modified diet.

For comparative nutritional information on all of our safflower oil products write:

Consumer Products Division, PVO

International Inc., World Trade Center,

San Francisco,
California
94111.



LIMITED WARRANTY TO CONSUMER
★ Good Housekeeping
★ PROMISES
★ REPLACEMENT OR REFUND IF DEFECTIVE

Saffola®

Highest in beneficial polyunsaturates.

*The Society for Continuing Medical Education
proudly announces:*

**AN INTERNATIONAL SYMPOSIUM ON
GYNECOLOGIC ONCOLOGY**

**LOEWS MONTE-CARLO HOTEL
May 31 through June 3, 1978**

**Groups departing from the United States will leave on
May 27th and are scheduled to return on June 5th.**

Program Chairman:
DR. DONALD WOODRUFF
The Johns Hopkins Medical Institutions

Program European Coordinator:
PROFESSOR DR. PER KOLSTAD
Director of Norwegian Radium Hospital

**This course has been approved for a total of 40 Cognates by The
American College of Obstetricians and Gynecologists. This program
is also approved for 24 Category 1 continuing medical education
credits.**

**Group rates for land arrangements have been made by the society
in an extremely attractive package.**

**Registration Fee: Members: \$125.00
Non-Members: \$175.00**

Air travel arrangements are via Sabena Belgian World Airlines

ATTENDANCE LIMITED

For information, please contact:

DR. ERWIN WITKIN
Society for Continuing Medical Education, Inc.
6609 Reisterstown Road, Suite 208
Baltimore, Maryland 21215

**Announcing the
41st Annual New Orleans
Medical Assembly**

**March 31 - April 4, 1978
The Fairmont—New Orleans**

THE HIGH RISK PATIENT

Accreditation: AMA Category 1, Physicians'
Recognition Award, American Academy of
Family Practice, American College of Emer-
gency Medicine, Category 1.

**ADOLPH A. FLORES, JR., M.D., President; OLIVER H.
DABEZIES, JR., M.D., F.A.C.S., Director of Program.**

Registration Fee: \$200 non-member physi-
cians; \$100 military; \$100 registered
nurses; Complimentary: students, resi-
dents, interns and Fellows.

Write or Phone: New Orleans Graduate Medi-
cal Assembly, Room 1538, Tulane Medical
Center, 1430 Tulane Ave., New Orleans,
LA 70112. Call (504) 525-9930.

• Presidential Reception • Exhibitors' Cham-
pagne Reception • Mississippi River Cruise
aboard SS Natchez • Superdome Tour, lunch-
eon & fashion show for wives & guests.

Spring Fiesta, March 31, 1978.
"A Night in Old New Orleans", Jackson Square.

ANNUAL SOUTH BAY HOSPITAL MEDICAL SYMPOSIUM CRITICAL CARE — 1978

**Friday, March 3, 1978
8:00 AM — 5:00 PM**

**HOLIDAY INN
21333 Hawthorne Blvd.
Torrance, California 90503**

**Saturday, March 4, 1978
9:00 AM — NOON**

GUEST FACULTY

Bernard Lown, M.D.

Michael Criley, M.D.

Richard Glassock, M.D.

Lucien Guze, M.D.

Paul Selecky, M.D.

H. J. C. Swan, M.D.

Max Harry Weil, M.D.

**Credit: 12 hours Category I CME Credit
in association with Harbor General Hospital.**


Registration Fees

Physicians \$75.00
RN's \$35.00
(Includes luncheon)

Mail to:

South Bay Hospital
Medical Education Dept.
514 N. Prospect Ave.
Redondo Beach, CA 90277

Brochure on Request



Just 2
days prior
to the Ameri-
can College of
Cardiology An-
nual Meeting in
Anaheim ...
Freeway-close in the
sunny, South Bay Beach
area.

VALIUM[®] (diazepam)^{IV} can effectively relieve anxiety and its somatic symptoms.

Initial calming in hours

Your anxious patient will be reassured by the prompt action of Valium.
It's immediate, tangible proof that the medication is working.

Significant improvement in days

Noticeable improvement of anxiety symptoms is usually evident within
the first few days of therapy.

Patient response you know, want and trust

Valium offers clinical effectiveness and a
wide margin of safety, which makes it a prudent choice for treating
psychic tension and anxiety.

Before prescribing, please consult complete product information, a summary of which follows:


Indications: Tension and anxiety states; somatic complaints which are concomitants of emotional factors; psychoneurotic states manifested by tension, anxiety, apprehension, fatigue, depressive symptoms or agitation; symptomatic relief of acute agitation, tremor, delirium tremens and hallucinosis due to acute alcohol withdrawal; adjunctively in skeletal muscle spasm due to reflex spasm to local pathology; spasticity caused by upper motor neuron disorders; athetosis; stiff-man syndrome; convulsive disorders (not for sole therapy).

Contraindicated: Known hypersensitivity to the drug. Children under 6 months of age. Acute narrow angle glaucoma; may be used in patients with open angle glaucoma who are receiving appropriate therapy.

Warnings: Not of value in psychotic patients. Caution against hazardous occupations requiring complete mental alertness. When used adjunctively in convulsive disorders, possibility of increase in frequency and/or severity of grand mal seizures may require increased dosage of standard anticonvulsant medication; abrupt withdrawal may be associated with temporary increase in frequency and/or severity of seizures. Advise against simultaneous ingestion of alcohol and other CNS depressants. Withdrawal symptoms (similar to those with barbiturates and alcohol) have occurred following abrupt discontinuance (convulsions, tremor, abdominal and muscle cramps, vomiting and sweating). Keep addiction-prone individuals under careful surveillance because of their predisposition to habituation and dependence.

Usage in Pregnancy: Use of minor tranquilizers during first trimester should almost always be avoided because of increased risk of congenital malformations as suggested in several studies. Consider possibility of pregnancy when instituting therapy; advise patients to discuss therapy if they intend to or do become pregnant.

Precautions: If combined with other psychotropics or anticonvulsants, consider carefully pharmacology of agents employed; drugs



such as phenothiazines, narcotics, barbiturates, MAO inhibitors and other antidepressants may potentiate its action. Usual precautions indicated in patients severely depressed, or with latent depression, or with suicidal tendencies. Observe usual precautions in impaired renal or hepatic function. Limit dosage to smallest effective amount in elderly and debilitated to preclude ataxia or oversedation.

Side Effects: Drowsiness, confusion, diplopia, hypotension, changes in libido, nausea, fatigue, depression, dysarthria, jaundice, skin rash, ataxia, constipation, headache, incontinence, changes in salivation, slurred speech, tremor, vertigo, urinary retention, blurred vision. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances, stimulation have been reported; should these occur, discontinue drug. Isolated reports of neutropenia, jaundice; periodic blood counts and liver function tests advisable during long-term therapy.

Dosage: Individualize for maximum beneficial effect.


Adults: Tension, anxiety and psychoneurotic states, 2 to 10 mg b.i.d. to q.i.d.; alcoholism, 10 mg t.i.d. or q.i.d. in first 24 hours, then 5 mg t.i.d. or q.i.d. as needed; adjunctively in skeletal muscle spasm, 2 to 10 mg t.i.d. or q.i.d.;

adjunctively in convulsive disorders, 2 to 10 mg b.i.d. to q.i.d. **Geriatric or debilitated patients:** 2 to 2½ mg, 1 or 2 times daily initially, increasing as needed and tolerated. (See Precautions.) **Children:** 1 to 2½ mg t.i.d. or q.i.d. initially, increasing as needed and tolerated (not for use under 6 months).

Supplied: Valium[®] (diazepam) Tablets, 2 mg, 5 mg and 10 mg—bottles of 100 and 500; Tel-E-Dose[®] packages of 100, available in trays of 4 reverse-numbered boxes of 25, and in boxes containing 10 strips of 10; Prescription Paks of 50, available singly and in trays of 10.



Roche Laboratories
Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110



This is a source
of somatic symptoms
of anxiety.



And this is a source of relief...

VALIUM[®]
(diazepam)^{IV}

2-mg, 5-mg, 10-mg scored tablets



Please see the preceding page for a summary of product information.

Artist's symbolic conception of a CNS reaction to excessive stress in the overanxious patient, resulting in somatic symptoms of the gastrointestinal and cardiovascular systems. Special photographic lighting techniques were applied to a model of the brain.